

Circumvent® PCV M



Porcine Circovirus Vaccine, Type 2, Killed Baculovirus Vector
Mycoplasma Hyopneumoniae Bacterin

An aid in the prevention of viremia and an aid in the reduction of virus shedding caused by porcine circovirus type 2, and an aid in the control of pneumonia caused by Mycoplasma hyopneumoniae



The new gold standard for PCV2 and M. hyo protection

- Combines Gold Standard PCV2 and M. hyo protection
- Ready to use combination assurance
- Ready to use combination convenience
- Proven prevention of chronic PCV2 infection (viremia)
- Proven reduction of PCV2 shedding
- Effective, long-lasting 2-dose protection
- No requirement to use within 4 hours or discard thereafter
- No risk of contamination due to mixing
- No additional labor for mixing or administration
- No second bottle to dispose of cutting waste by half
- Sustained protection to market without fear of late breaks
- Allows for maximum performance and less variation in finishing

Circumvent® PCV M

Aids in the prevention of viremia, and as an aid in the reduction of virus shedding caused by porcine circovirus type 2, and as an aid in the control of pneumonia caused by *Mycoplasma hyopneumoniae*.

Circumvent® PCV M

Porcine Circovirus Vaccine, Type 2, Killed Baculovirus Vector, *Mycoplasma Hyopneumoniae* Bacterin

INDICATIONS AND CONTENTS:

For use in healthy swine, 3 weeks of age or older, as an aid in the prevention of viremia, and as an aid in the reduction of virus shedding caused by porcine circovirus type 2, and as an aid in the control of pneumonia caused by *Mycoplasma hyopneumoniae*.

DOSAGE AND DIRECTIONS: Allow the vaccine to reach an ambient temperature of 16°- 27° C (60°- 80° F). Shake well and administer a 2.0 mL dose intramuscularly to healthy swine, 3 weeks of age or older, followed by a second 2.0 mL dose 3 weeks later. Duration of immunity is under development.

CAUTIONS: Store refrigerated at 2°-7°C (35°-45°F). Do not freeze. Use immediately after opening; do not save partial contents. Burn the container and all unused product. Use only in healthy swine. Do not vaccinate within 21 days of slaughter. Caution should be used to avoid injecting your own fingers, hands, or other body parts. Accidental injection can cause serious local reactions. Contact a physician immediately.

Please note that local reactions may occur at the injection sites in vaccinates, and most of these reactions resolve in 16 to 23 days following vaccination. Some systemic reactions may be seen following vaccination. If allergic reaction occurs, treat with epinephrine.

PRESERVATIVES: Gentamicin and thimerosal are added as preservatives.

SUPPLIED:	Code: 046494	500 mL 250 Doses
	Code: 029726	100 mL 50 Doses

FOR VETERINARY USE ONLY

READ AND FOLLOW LABEL DIRECTIONS

U.S. Veterinary License No. 165A



Intervet/Schering-Plough Animal Health • 556 Morris Ave. • Summit, NJ 07901 • www.intervetusa.com • 800.521.5767

Circumvent and Pig Head logo are the property of Intervet International B.V. or affiliated companies or licensors and are protected by copyrights, trademark and other intellectual property laws. ©2011 Intervet International B.V. All rights reserved. • Part No. SW-CIRM-PB • MAG 1/11