

Porcine Circovirus Vaccine, Type 2, Killed Baculovirus Vector, Mycoplasma Hyopneumoniae Bacterin



Circumvent[®] PCV M

250 doses 500 mL Code 046494



Intervet
Schering-Plough Animal Health

INTERVET INC., Omaha, NE 68103, USA
www.intervetusa.com 1-800-521-5767
U.S. Veterinary License No. 165A

For use in healthy swine, 3 weeks of age or older, as an aid in the prevention of viremia, and as an aid in the reduction of virus shedding caused by porcine circovirus type 2, and as an aid in the control of pneumonia caused by *Mycoplasma hyopneumoniae*.

DOSE: Allow the vaccine to reach an ambient temperature of 16°- 27° C (60°- 80° F). Shake well and administer a 2.0 mL dose intramuscularly to healthy swine, 3 weeks of age or older, followed by a second 2.0 mL dose 3 weeks later. Duration of immunity is under development.


CAUTIONS: Store refrigerated at 35°- 45°F (2°- 7°C). Do not freeze. Use immediately after opening; do not save partial contents. Burn the container and all unused product. Use only in healthy swine. Do not vaccinate within 21 days of slaughter. Caution should be used to avoid injecting your own fingers, hands, or other body parts. Accidental injection can cause serious local reactions. Contact a physician immediately.

Please note that local reactions may occur at the injection sites in vaccinates, and most of these reactions resolve in 16 to 23 days following vaccination. Some systemic reactions may be seen following vaccination. If allergic reaction occurs, treat with epinephrine.

PRESERVATIVES: Gentamicin and thimerosal are added as preservatives.

FOR VETERINARY USE ONLY

905925002



Serial No:

Exp. Date: